

**ON ALMOST STABLE CMC HYPERSURFACES IN
MANIFOLDS OF BOUNDED SECTIONAL
CURVATURE**

Julien Roth, Abhitosh Upadhyay

► **To cite this version:**

Julien Roth, Abhitosh Upadhyay. ON ALMOST STABLE CMC HYPERSURFACES IN MANIFOLDS OF BOUNDED SECTIONAL CURVATURE. 2019. hal-01995976

HAL Id: hal-01995976

<https://hal-upec-upem.archives-ouvertes.fr/hal-01995976>

Preprint submitted on 28 Jan 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ON ALMOST STABLE CMC HYPERSURFACES IN MANIFOLDS OF BOUNDED SECTIONAL CURVATURE

JULIEN ROTH AND ABHITOSH UPADHYAY

ABSTRACT. We show that almost stable constant mean curvature hypersurfaces contained in a sufficiently small ball of a manifold of bounded sectional curvature are close to geodesic spheres.

1. INTRODUCTION AND STATEMENT OF THE RESULT

Let $\phi : (M^n, g) \rightarrow (N^{n+1}, \bar{g})$ be an isometric immersion of an oriented closed n -dimensional Riemannian manifold M in a $(n + 1)$ -dimensional oriented manifold (N, \bar{g}) . We assume that M is oriented by the global unit normal field ν so that ν is compatible with the orientations of M and N . We will denote by B the second fundamental form of ϕ and its mean curvature by H . Let $F : (-\varepsilon, \varepsilon) \times M \rightarrow N$ be a variation of ϕ so that $F(0, \cdot) = \phi$. The balance volume associated with the variation F is the function $V : (-\varepsilon, \varepsilon) \rightarrow \mathbb{R}$ defined by

$$\int_{[0,t] \times M} F^* dv_{\bar{g}}$$

where $dv_{\bar{g}}$ is the volume element associated to the metric \bar{g} on N . We will denote simply by dv the volume element of g . It is a classical fact that

$$V'(0) = \int_M f dv$$

where $f(x) = \langle \frac{\partial F}{\partial t}(0, x), \nu \rangle$. Moreover the area function $A(t) = \int_M dv_{F_t^* h}$ satisfies

$$A'(0) = -n \int_M H f dv.$$

We say that F is preserving the volume if $V(t) = V(0)$ in a neighbourhood of 0 and in this case we have $\int_M f dv = 0$. Conversely, for all smooth functions f so that

$\int_M f dv = 0$, there exists a preserving volume variation so that $f = \langle \frac{\partial F}{\partial t}(0, x), \nu \rangle$.

The following assertions are equivalent

- (i) The immersion ϕ is a critical point of the area (i.e. $A'(0) = 0$) for all variations with preserving volume.
- (ii) $\int_M H f dv = 0$ for any smooth function so that $\int_M f dv = 0$.

2010 *Mathematics Subject Classification.* 53C24, 53C42.

Key words and phrases. Stable constant mean curvature, Hypersurface, Geodesic sphere.

- (iii) There exists a constant H_0 so that $A'(0) + nH_0V'(0) = 0$ for any variation.
- (iv) ϕ is of constant mean curvature H_0 .

An immersion with constant mean curvature H_0 will be said stable if $A''(0) \geq 0$ for all variations preserving the volume. Now we consider the function $\mathcal{J}(t)$ defined by

$$\mathcal{J}(t) = A(t) + nH_0V(t).$$

Then, $\mathcal{J}''(0)$ depends only on f and we have

$$\begin{aligned} \mathcal{J}''(0) &= \int_M |df|^2 dv - \int_M (Ric^N(\nu, \nu) + |B|^2) f^2 dv \\ &= \int_M f \Delta f - \int_M (Ric^N(\nu, \nu) + |B|^2) f^2 dv \\ &= \int_M f J f dv, \end{aligned}$$

where Ric^N is the Ricci curvature of N with respect to the metric \bar{g} and J is the so-called Jacobi operator defined by $Jf = \Delta f - (Ric^N(\nu, \nu) + |B|^2)f$. It is well known that ϕ is a stable constant mean curvature immersion if and only if $\mathcal{J}''(0) \geq 0$ for any smooth function f so that $\int_M f dv = 0$ or equivalently if J is a non-negative operator.

Barbosa and Do Carmo [1] proved that the only stable closed CMC hypersurfaces of the Euclidean space are the round spheres. This result was extended later by Barbosa, Do Carmo and Eschenburg [2] for spheres and hyperbolic spaces. In [4], Grosjean with the first author considered the stability of CMC hypersurfaces in Riemannian manifolds with (non-constant) bounded sectional curvature. After proving a pinching result for the first eigenvalue of the Laplacian, they were able to show that a closed stable CMC hypersurface of a Riemannian manifold with bounded sectional curvature and contained in a geodesic ball of sufficiently small radius is *close* to a geodesic sphere. Namely, *close* means here diffeomorphic and almost-isometric to a geodesic sphere of appropriate radius (depending upon the mean curvature).

The aim of this short note is to show that the assumption of being stable can be relaxed to almost stable in the result of [4]. By *almost stable*, we mean that the Jacobi operator J is not supposed to be non-negative but greater than some small negative constant

$$(1) \quad \int_M f J f dv \geq -n\varepsilon \int_M h^2 f^2 dv,$$

for any smooth function f so that $\int_M f dv = 0$, where $h = \sqrt{\|H\|_\infty^2 + \delta}$. Note that h^2 appears in the right hand side of the almost stability condition for homogeneity reasons.

From now on, we assume that the sectional curvature of (N, \bar{g}) satisfies $\mu \leq Sect_N \leq \delta$, for $\mu \leq \delta$ two real constants. Before stating the main result of this

note, we introduce the following function

$$s_\delta(r) = \begin{cases} \frac{1}{\sqrt{\delta}} \sin \sqrt{\delta} r & \text{if } \delta > 0 \\ r & \text{if } \delta = 0 \\ \frac{1}{\sqrt{|\delta|}} \sinh \sqrt{|\delta|} r & \text{if } \delta < 0, \end{cases}$$

as well as the following set: $\mathcal{H}_V(n, N)$ is the space of all Riemannian compact, connected and oriented n -dimensional Riemannian manifolds without boundary isometrically immersed into (N^{n+1}, \bar{g}) which satisfy the following hypothesis on the volume : $V(M) \leq \frac{c\omega_n}{\delta^{n/2}}$ if $\delta > 0$ and $V(M) \leq c\omega_n i(N)^n$ if $\delta \leq 0$ for some constant c . For more convenience we take $1/\sqrt{\delta} = +\infty$ if $\delta \leq 0$. This condition on the volume is required to apply the result of [4] (see Theorem 2.1 below) about the pinching of the first eigenvalue of the Laplacian. This condition comes from the extrinsic Sobolev inequality of Hoffman and Spruck [6] which is used in the proof of that pinching result.

The main result of this note is the following:

Theorem 1.1. *Let (N^{n+1}, \bar{g}) be a $(n+1)$ -dimensional Riemannian manifold whose sectional curvature $Sect_N$ satisfies $\mu \leq Sect_N \leq \delta$ and $i(N) \geq \frac{\pi}{\sqrt{\delta}}$ if $\delta > 0$ and let $M \in \mathcal{H}_V(n, N)$. Let us assume that $\phi(M)$ lies in a convex ball of radius $\min\left(\frac{\pi}{8\sqrt{\delta}}, \frac{i(N)}{2}\right)$. Let $\varepsilon < \frac{1}{12}$, $q > n$ and $A > 0$. Let assume that $V(M)^{1/n} \|B\|_q \leq A$ for $\delta \geq 0$ (resp. $\max(\frac{H}{h}, V(M)^{1/n} \|B\|_q) \leq A$ for $\delta < 0$). Then there exist positive constants $\alpha := \alpha(q, n)$, $K := K(n, q, A)$ and $R_0(\delta, \mu, \varepsilon)$ such that if ϕ is of constant mean curvature H and almost stable in the sense of (1), $\varepsilon^\alpha < 1/K$ and $\phi(M)$ is contained in a convex ball of radius $R_0(\delta, \mu, \varepsilon)$ then M is diffeomorphic and $K\varepsilon^\alpha$ -quasi-isometric to $S(p, s_\delta^{-1}(\frac{1}{h}))$, that is there exists a diffeomorphism F from M into $S(p, s_\delta^{-1}(\frac{1}{h}))$ so that*

$$||dF_x(u)|^2 - 1| \leq K\varepsilon^\alpha$$

for any $x \in M$, $u \in T_x M$ and $|u| = 1$.

2. PROOF OF THE RESULT

Let $M \in \mathcal{H}_V(n, N)$ and denote by ϕ the isometric immersion of (M, g) into (N, h) . Moreover, let us assume that ϕ has constant mean curvature and is almost stable in the sense of (1) for some positive ε . Let f be an eigenfunction associated with the first eigenvalue $\lambda_1(M)$ of the Laplacian on (M, g) . Since $\int_M f dv = 0$ then condition (1) gives

$$\lambda_1(M) \int_M f^2 dv - \int_M (Ric^N(\nu, \nu) + nH^2 + |\tau|^2) f^2 dv \geq -\varepsilon n h^2 \int_M f^2 dv$$

where τ is traceless part of the second fundamental form B . Since $\mu \leq \text{Sect}^N$, we deduce that

$$\begin{aligned} \lambda_1(M) &\geq n(H^2 + \mu - \varepsilon h^2) \\ &\geq nh^2 + n(\mu - \delta - \varepsilon h^2) \\ &\geq nh^2 \left(1 + \frac{\mu - \delta - \varepsilon h^2}{h^2} \right) \end{aligned}$$

and so

$$nh^2 \leq \lambda_1(M) \left(1 + \frac{1}{\frac{\delta - \mu + \varepsilon h^2}{h^2} - 1} \right).$$

Now, we set $R_1(\delta, \mu, \varepsilon) = s_\delta^{-1} \left(\sqrt{\frac{\frac{1}{2} - \varepsilon}{(\delta - \mu) \left(\frac{1}{2\varepsilon} + 1 \right)}} \right)$. We recall that the extrinsic radius of M is defined as the radius of the smallest ball containing $\phi(M)$. We have the following well-known lower bound for the extrinsic radius $R(M)$ (see [3]) for instance)

$$s_\delta(R(M)) \geq \frac{1}{h},$$

for compact hypersurfaces of a Riemannian manifold of sectional curvature bounded from above by δ . Now, if we assume that $\phi(M)$ is contained in a ball of radius R_1 , then since s_δ is an increasing function, we get

$$h^2 \geq \frac{1}{s_\delta^2(R(M))} \geq \frac{1}{s_\delta^2(R_1)} = \frac{(\delta - \mu) \left(\frac{1}{\varepsilon} + 1 \right)}{\frac{1}{2} - \varepsilon}.$$

From this, we deduce easily that

$$\frac{1}{\frac{\delta - \mu + \varepsilon h^2}{h^2} - 1} \leq 2\varepsilon$$

and so

$$nh^2 \leq \lambda_1(M)(1 + 2\varepsilon),$$

that is, $(\Lambda_{2\varepsilon})$ holds, where we denote by (Λ_η) the following pinching condition

$$(\Lambda_\eta) \quad nh^2 \leq \lambda_1(M)(1 + \eta)$$

associated with the following upper bound for the first eigenvalue of the Laplacian proved by Heintze [5]

$$\lambda_1(M) \leq nh^2.$$

Now, we recall the following theorem proved by the first author and Grosjean in [4].

Theorem 2.1 ([4]). *Let (N^{n+1}, \bar{g}) be a $n+1$ -dimensional Riemannian manifold whose sectional curvature Sect_N satisfies $\mu \leq \text{Sect}_N \leq \delta$ and $i(N) \geq \frac{\pi}{\sqrt{\delta}}$ if $\delta > 0$. Let $M \in \mathcal{H}_V(n, N)$. Let us assume that $\phi(M)$ lies in a convex ball of radius $\min\left(\frac{\pi}{8\sqrt{\delta}}, \frac{i(N)}{2}\right)$. Let p_0 be the center of mass of M . Let $\eta < 1/6$, $q > n$ and $A > 0$. Let us assume that $\max(V(M)^{1/n}\|H\|_\infty, V(M)^{1/n}\|B\|_q) \leq A$ for $\delta \geq 0$ (resp. $\max(V(M)^{1/n}\|H\|_\infty, \frac{\|H\|_\infty}{h}, V(M)^{1/n}\|B\|_q) \leq A$ for $\delta < 0$). Then there exist positive constants $C := C(n, q, A)$ and $\alpha := \alpha(q, n)$ such that if*

(Λ_η) holds, $\eta^\alpha < 1/C$ and $\phi(M)$ is contained in the ball $B\left(p_0, s_\delta^{-1}\left(\sqrt{\frac{\eta}{\delta-\mu}}\right)\right)$ then M is diffeomorphic and $C\eta^\alpha$ -quasi-isometric to $S(p, s_\delta^{-1}(\frac{1}{h}))$.

Let $\varepsilon < \frac{1}{12}$ and $\eta = 2\varepsilon < \frac{1}{6}$. We set

$$R_0(\delta, \mu, \varepsilon) = \min \left\{ s_\delta^{-1} \left(\sqrt{\frac{2\varepsilon}{\delta-\mu}} \right), R_1(\delta, \mu, \varepsilon) \right\}$$

and $K(n, q, A) = 2^{\alpha(n, q)} C(n, q, A)$, where α , C and R are the constants given by Theorem 2.1 and R_1 is defined in the beginning of the proof.

If we assume that $\phi(M)$ is contained in a ball of radius $R_0(\delta, \mu, \varepsilon)$, by the definition of $R_0(\delta, \mu, \varepsilon)$, we have $R_0(\delta, \mu, \varepsilon) \leq R_1(\delta, \mu, \varepsilon)$ and so we get from the above computation that $(\Lambda_{2\varepsilon})$ holds. Moreover, we also get that $\phi(M)$ is contained in a ball of radius $s_\delta^{-1}\left(\sqrt{\frac{2\varepsilon}{\delta-\mu}}\right)$. In addition, if we assume that $\varepsilon^\alpha < 1/K$, then, from the definition of K , we get that $(2\varepsilon)^\alpha < 1/C$ and Theorem 2.1 (applied with $\eta = 2\varepsilon$) gives that M is diffeomorphic and $C\eta^\alpha$ -quasi-isometric to $S(p, s_\delta^{-1}(\frac{1}{h}))$. Since $C\eta^\alpha = K\varepsilon^\alpha$, by the definition of K , Theorem 1.1 is proven. \square

Remark 2.2. *In the case of the Euclidean space, the result of Theorem 1.1 was proven in [7] and in this case, there is no assumption of being contained in a small ball. Theorem 1.1 gives an analogue of the result of [7] for the hyperbolic space as well as the half-sphere. Indeed, if $\mu = \delta$, then $R_0(\delta, \mu, \varepsilon) = +\infty$ since both $s_\delta^{-1}\left(\sqrt{\frac{2\varepsilon}{\delta-\mu}}\right)$ and $R_1(\delta, \mu, \varepsilon)$ are $+\infty$.*

Acknowledgements

Second author gratefully thanks the support of National Post-doctoral Fellowship (**File no. PDF/2017/001165**) from Science and Engineering Research Board, Government of India.

REFERENCES

- [1] J.L. BARBOSA & M.P. DO CARMO Stability of Hypersurfaces with Constant Mean Curvature, *Math. Z.*, **185**, No. 3, (1984), 339-353.
- [2] J.L. Barbosa, M.P. do Carmo & J. Eschenburg, *Stability of hypersurfaces of constant mean curvature in Riemannian manifolds*, *Math. Z.* **197** (1) (1988), 123-138.
- [3] C. Baikoussis & T. Koufogiorgos, *The diameter of an immersed Riemannian manifold with bounded mean curvature*, *J. Austral. Soc.*, (Series A), **31** (1981), 189-192.
- [4] J.F. Grosjean & J. Roth, *Eigenvalue Pinching and Application to the Stability and the Almost Umbilicity of Hypersurfaces*, *Math. Z.* **271**(1) (2012), 469-488.
- [5] E. Heintze, *Extrinsic upper bound for λ_1* , *Math. Ann.*, **280**, (1988), 389-402.
- [6] D. Hoffman & J. Spruck, *Sobolev and isoperimetric inequalities for Riemannian submanifolds*, *Comm. Pure and Appl. Math.*, **27** (1974), 715-727.
- [7] J. Roth & J. Scheuer, *Pinching of the first eigenvalue for second order operators on hypersurfaces of the Euclidean space*, *Ann. Glob. Anal. Geom.* **51**(3) (2017), 287-304.

(J. ROTH) LABORATOIRE D'ANALYSE ET DE MATHÉMATIQUES APPLIQUÉES, UPEM-UPEC, CNRS,
F-77454 MARNE-LA-VALLÉE

E-mail address: `julien.roth@u-pem.fr`

(A. UPADHYAY) INDIAN INSTITUTE OF SCIENCE, DEPARTMENT OF MATHEMATICS, BANGALORE,
560012, INDIA

E-mail address: `abhi.basti.ipu@gmail.com`, `abhitoshu@iisc.ac.in`